

RESEARCH

Open Access



Increased ex vivo cell death of central memory CD4 T cells in treated HIV infected individuals with unsatisfactory immune recovery

Marta Massanella^{1,4}, Elisabet Gómez-Mora¹, Jorge Carrillo¹, Marta Curriu¹, Dan Ouchi¹, Jordi Puig², Eugènia Negro^{2,3}, Cecilia Cabrera¹, Bonaventura Clotet^{1,2,3} and Julià Blanco^{1,3*}

Abstract

Background: High levels of ex vivo CD4 T-cell death and the accumulation of highly differentiated and/or immunosenescent T cells have been associated with poor CD4 T-cell recovery in treated HIV-infected individuals. However, the relationship between cell death and T-cell differentiation is still unclear.

Methods: We have analyzed cell death, immunosenescence and differentiation parameters in HAART-treated subjects (VL <50 copies/mL for more than 2 years) with CD4 T-cell count <350 cells/ μ L (immunodiscordant, n = 23) or >400 cells/ μ L (immunoconcordant, n = 33). We included 11 healthy individuals as reference.

Results: As expected, suboptimal CD4 T-cell recovery was associated with low frequencies of naïve cells, high frequencies of transitional and effector memory cells and a subsequent low ratio of central/transitional memory cells in the CD4 compartment. These alterations correlated with spontaneous CD4 T-cell death. A deeper analysis of cell death in CD4 T-cell subsets showed increased cell death in memory cells of immunodiscordant individuals, mainly affecting central memory cells. Immunosenescence was also higher in immunodiscordant individuals albeit unrelated to cell death. The CD8 compartment was similar in both HIV-infected groups, except for an underrepresentation of naïve cells in immunodiscordant individuals.

Conclusion: Immunodiscordant individuals show alterations in memory CD4 T-cell differentiation associated with a short ex vivo lifespan of central memory cells and an in vivo low central/transitional memory cell ratio. These alterations may contribute to poor CD4 T-cell repopulation.

Keywords: Immunodiscordant, Cell death, T-cell subsets, Immunosenescence, HAART, cART

Background

HIV-1 infection leads to major perturbations of immune regulatory mechanisms [1] as a consequence of virus-mediated depletion of CD4 T cells [2, 3] and increased immune activation and inflammation [1]. Abnormal proliferative responses, maturation and immunosenescence of T cells, affecting both CD4 and CD8 compartments

and covering acute, chronic and late stages of infection have been reported [4].

Highly active antiretroviral therapy (HAART) efficiently controls HIV-1 replication, but inflammatory status and immune activation remain abnormally high in treated individuals compared with uninfected individuals [1]. Furthermore, CD4 T-cell recovery shows a heterogeneous response [5, 6]; while some subjects reach CD4 T-cell counts comparable to healthy individuals, others fail to increase CD4 T-cell counts even after long-term (>5 years) virologically effective treatment [5, 7]. This immunodiscordant response has been associated with

*Correspondence: jblanco@irsicaixa.es

¹ Institut de Recerca de la Sida IrsiCaixa-HIVACAT, Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Universitat Autònoma de Barcelona, 08916 Badalona, Spain

Full list of author information is available at the end of the article

immunological damage induced by HIV-1 replication measured by nadir CD4 T-cell counts [7, 8] and specifically with the destruction of thymic tissue and the consequent lack of production of new naïve cells [9, 10]. However, in the absence of new T cells, memory cells can repopulate the CD4 T-cell compartment in non-human primate models of HIV infection [11], suggesting that the damage of memory cells may also contribute to blunted immunological recoveries.

Increased replicative senescence, measured by the senescence marker CD57 [12], have been related with immune recovery [13] and could affect CD4 T cells in both naïve and memory subsets [14]. Apart from replicative senescence, skewed maturation of memory CD4 T cells [15] and higher turnover and sensitivity to spontaneous cell death [9, 15] may contribute to a lower proliferative potential of memory cells leading to incomplete immune recovery under HAART. However, it is still unclear how these different alterations of memory cells are linked and how are they associated with thymic damage or senescence. Moreover, the specific role of central memory CD4 T cells, a major player of HIV pathogenesis [16], in immune recovery remains poorly defined.

Considering the role of CD4 T-cell death in immunological recovery on HAART [6], we sought to analyze its relationship with altered maturation/immunosenescence profiles of memory cells. Therefore, we designed a study to compare these parameters in subjects with low and high CD4 T-cell recovery after long-term successful HAART. An aberrant maturation phenotype defined by a reduced central/transitional memory CD4 T-cell ratio, but not immunosenescence, strongly correlated with poor CD4 T-cell recovery and seems to be associated with high cell death levels in intermediate differentiation stages of memory cells.

Methods

Samples

A cross-sectional study was performed to analyze the level of immunosenescence and differentiation of T cells in HAART-treated subjects. The Institutional Review Board of the Hospital Germans Trias i Pujol approved the study (EO code: EO-07-024) and all individuals provided written informed consent. Subjects that participated in the APOP-V⁺I⁻ study [8, 17] were screened for the current study: 56 individuals with at least two frozen aliquots of PBMCs were selected. All subjects had confirmed diagnosis of HIV infection, continuous HAART with plasma viral load <50 copies/mL for at least the last 2 years (minimum of 4 determinations during this period) and good treatment adherence. Exclusion criteria were: chemotherapy or interferon/ribavirin treatment

and history of opportunistic infections during the last 2 years.

Patients were classified as Concordant (favorable virologic and immunologic response) when CD4 T-cell counts were above 400 cells/ μ L. Conversely, Discordant patients (favorable virologic but unsatisfactory immunologic response) showed CD4 T cells <350 cells/ μ L [8, 17]. Clinical and demographic data were collected from medical records. For comparative purposes, frozen PBMC from a control group of HIV uninfected individuals ($n = 11$; EO-10-007, Institutional Review Board of the Hospital Germans Trias i Pujol) were also analyzed.

CD4 and CD8 T-cell immunophenotype

Frozen PBMCs were thawed at 37°C, washed twice in RPMI/20% of fetal bovine serum (FBS) and incubated for 1 h at 37°C before staining with a previously described eight-color panel [18] including CD3-APC-Cy7, CD4-PerCP-Cy5.5, CD8-V500, CD57-FITC, CD27-APC, CD28-PE, CCR7-PE-Cy7 and CD45RA-V450 (BD Biosciences). Cell viability was evaluated by Propidium Iodide (PI, Sigma-Aldrich) staining in parallel tubes, the median [IQR] of PI⁺ cells was 2.1% [1.5–4.0]. Lymphocyte gate was defined by morphological parameters excluding most dead cells (less than 0.1% of PI⁺ cells were found in the living lymphocyte gate). At least 300,000 lymphocytes were acquired in a LSRII cytometer (BD Biosciences) and analyzed with FlowJo software (Tree Star). Gating strategy is described in Additional file 1: Figure S1. Replicative senescence was defined by CD57⁺CD28⁻ cells. T-cell subsets were first analyzed for CD27 and CD28 expression to identify Terminally Differentiated cells (T_{TD}, CD28⁻CD27⁻) and Effector Memory cells (TEM, CD27⁻CD28⁺ for CD4 T cells or CD27⁺CD28⁻ for CD8 T cells). Double positive CD27⁺CD28⁺ cells were sub-analyzed for CD45RA and CCR7 expression allowing for the definition of naïve (T_N, CD45RA⁺CCR7⁺), central memory (T_{CM}, CD45RA⁻CCR7⁺) and transitional memory cells (T_{TM}, CD45RA⁻CCR7⁻). For each subset the expression of CD57 was analyzed (Additional file 1: Figure S1).

Immune activation, absolute lymphocyte counts (used to calculate absolute counts of each subset) and thymic production (CD45RA⁺CD31⁺CD4 T cells) were previously assessed in fresh blood samples in both CD4 and CD8 T cells [8, 17].

Ex vivo culture of whole PBMC and sorted cells

Data on spontaneous CD4 and CD8 T-cell death (including necrosis, intrinsic and extrinsic apoptosis) have been previously reported and were obtained after culture of freshly obtained PBMC for 24 h [17, 19]. Furthermore,

in a subgroup of patients ($n = 10$) freshly obtained PBMC were stained with CD3-APC-Cy7, CD4-PerCP-Cy5.5, CD8-V500, CD27-APC, CCR7-PE-Cy7 and CD45RA-V450 as described above, washed in PBS and sorted immediately in a FACSAria (BD Biosciences) to obtain gated $CD3^+CD4^+CD8^-$ cells with the following phenotypes: T_N cells ($CD27^+CD45RA^+CCR7^+$), T_{CM} cells ($CD27^+CD45RA^-CCR7^+$), T_{TM} cells ($CD27^+CD45RA^-CCR7^-$) and a mixture of T_{EM} and T_{TD} cells ($T_{EM+TD} CD27^-$). These latter subsets were sorted as a single population due to technical limitations of sorting procedure (only 4 sorted populations could be obtained from a single sample). Sorted cells were cultured for 24 h and cell death was assayed using 40 nM DIOC(6) as described previously [19]. DIOC(6)^{low} cells, having lost mitochondrial membrane potential, were identified as dead cells.


Statistical analysis

Continuous variables were expressed as the median (interquartile range) and compared using Mann–Whitney U test, permutation test for unbalanced groups or Signed-Rank test (for paired analyses). Discrete variables were described as percentages and compared using the Fisher's exact test. Correlations were assessed by the Spearman's test. Multiple comparisons were adjusted for false discovery rate. Statistical analyses were performed using R software version 3.0.2 [20] with two-tailed significance levels of 5%.

Results

Subject characteristics

Subjects recruited in this study were stratified according to the level of recovery of CD4 T cells after suppressive HAART, as immunodiscordant or immunoconcordant (cutoff value 350 CD4 T cells/ μ L). The time course of CD4 T-cell recovery for each group is shown in Figure 1a. At the time of sample analysis, the median CD4 T-cell count was 220 and 798 cells/ μ L for immunodiscordant and immunoconcordant HIV-infected individuals respectively ($p < 0.0001$), reflecting the blunted dynamics of CD4 T-cell recovery of the former group. Absolute CD8 T-cell counts showed no differences between HIV-infected groups. Median length of HIV infection (from diagnosis) was 11.8 and 10.1 years ($p = ns$), while median time on treatment was 11.2 and 5.2 years ($p = ns$) for immunoconcordant or immunodiscordant subjects, respectively. Overall, immunodiscordant individuals showed the previously reported higher CD4 T-cell death and activation with significantly lower nadir CD4 T-cell counts [8, 17]. However, some immunoconcordant individuals showed also low nadir values. To evaluate the effect of nadir on immune recovery, immunoconcordant



individuals were further divided according to the median nadir CD4 T-cell value (cut-off value 250 CD4 T cells/ μ L) into two subgroups with low and high nadir values ($n = 17$ and $n = 16$, respectively). This allowed the direct comparison of immunoconcordant individuals with immunodiscordant subjects, avoiding the confounding

effect of nadir and enabling us to evaluate the impact of low nadir values in immunoconcordant individuals. No major differences were observed among immunoconcordant subgroups, except for a higher presence of PI-based treatments and HCV co-infection in the low nadir group (Table 1).

Analysis of the CD4 T-cell maturation

Absolute counts and frequency of different CD4 T-cell subsets were analyzed in immunodiscordant, immunoconcordant (low and high-nadir subgroups) and 11 uninfected individuals. The data show that lower CD4 T-cell counts in immunodiscordant subjects (Figure 1a) were the consequence of lower levels of T_{N^c} , T_{CM^c} , T_{TM} and

T_{EM} cells compared with immunoconcordant individuals, while the absolute numbers of TTD cells were similar in all groups (Figure 1b). Interestingly, immunoconcordant patients, irrespective of the nadir values showed similar counts of all subsets that were in turn comparable to uninfected controls except for the T_{EM} subset, suggesting a proper recovery of the CD4 T-cell subsets in these individuals (Figure 1b). The frequency of each subset in the CD4 T-cell compartment showed a significant underrepresentation of TN cells in immunodiscordant subjects (as compared to concordant or HIV-uninfected individuals) that was compensated by an overrepresentation of T_{TM} cells and a less evident but still significant increase in T_{EM} and T_{TD} cells (Figure 1c). Conversely, T_{CM} cells showed

Table 1 Main characteristics of the different groups

	Discordant (n = 23)	a	Concordant			b	Uninfected (n = 11)
			All (n = 33)	Low Nadir (n = 17)	High Nadir (n = 16)		
Age (years), Median [IQR]	48 [45–50]	ns	45 [38–49]	48 [42–52]	42 [37–45]	ns	38 [34–47]
Gender (% of male)	91	ns	85	76	94	ns	55
Time since HIV diagnosis (years), Median [IQR]	10.1 [4.1–20.4]	ns	11.8 [7.5–16.6]	12.2 [8.9–17.4]	11.6 [4.9–13.4]	ns	–
Time on HAART (years), Median [IQR]	5.2 [3.5–11.4]	ns	11.2 [7.4–12.6]	11.4 [8.8–12.6]	10.6 [3.7–12.8]	ns	–
Current HAART (% PI-based)	70	*	33	41	25	*	–
HCV coinfection (%)	35	ns	21	36	15	*	0
Ratio CD4/CD8, Median [IQR]	0.23 [0.17–0.33]	*	0.87 [0.60–1.11]	0.76 [0.54–0.87]	1.06 [0.86–1.1]	ns	1.64 [1.31–1.81]
CD4 T cell counts (cells/ μ L), Median [IQR]	220 [192–253]	*	798 [600–998]	703 [600–896]	881 [672–1,075]	ns	779 [629–1,072]
Nadir (cells/ μ L), Median [IQR]	64 [15–122]	*	239 [76–345]	76 [19–185]	351 [280–429]	*	–
CD4 T-cell gain (cell/ μ L/year HAART), Median [IQR]	27 [9–54]	*	53 [46–102]	53 [46–93]	53 [36–124]	ns	–
CD4 T cell (% of lymph), Median [IQR]	14 [10–17]	*	31 [27–39]	29 [27–33]	37 [29–40]	ns	42 [36–45]
CD4 T-cell death (%), Median [IQR]	9.3 [7.5–15.1]	*	4.6 [3.3–5.9]	4.9 [4.4–5.6]	4.3 [2.9–6.1]	ns	4.0 [3.2–4.7]
CD38 ⁺ CD45RA ⁻ (% of CD4 T cells), Median [IQR]	37 [29–41]	*	25.7 [20–32]	26.5 [20–32]	25.1 [20–31]	ns	–
HLA-DR ⁺ CD95 ⁺ (% of CD4 T cells), Median [IQR]	16 [7.7–21.6]	*	4.5 [3.7–6.7]	4.7 [4.2–6.3]	4.4 [3.2–7.1]	ns	2.0 [1.5–2.7]
CD8 T cell counts (cells/ μ L), Median [IQR]	940 [754–1,146]	ns	908 [771–1,239]	1,118 [855–1,380]	811 [637–1,121]	ns	459 [433–548]
CD8 T cell (% of lymph), Median [IQR]	56 [51–61]	*	38 [34–46]	39 [36–47]	36 [32–40]	ns	24 [23–27]
CD8 T cell death (%), Median [IQR]	7.1 [4.8–10.3]	ns	6.5 [5.0–12.4]	8.8 [6.1–11.0]	6.1 [4.8–14.9]	ns	3.9 [3.4–4.9]
CD38 ⁺ CD45RA ⁻ (% of CD8 T cells), Median [IQR]	31 [23–36]	*	23 [19–35]	25 [21–36]	21 [19–26]	ns	9 [5–13]
HLA-DR ⁺ CD95 ⁺ (% of CD8 T cells), Median [IQR]	12.6 [6.9–22.4]	ns	9.1 [5.6–13.0]	9.8 [5.4–13.6]	8.5 [5.6–13.0]	ns	–
sCD14 (μ g/mL), Median [IQR]	8.4 [7.7–10.2]	ns	8.8 [7.2–9.7]	9.2 [7.6–10.1]	8.0 [7.1–9.2]	ns	4.2 [3.9–4.6]

a Comparison of concordant and discordant subjects. * denotes $p < 0.05$; ns non significant (Mann–Withney U or Fisher exact test).

b Comparison of concordant subjects with low and high nadir. * denotes $p < 0.05$; ns non significant (Mann–Withney U or Fisher exact test).

similar values in all groups. Again, both subgroups of immunocordant subjects showed similar values of subset frequencies reaching the levels of HIV-uninfected controls (Figure 1c).

CD4 T-cell maturation and CD4 T-cell death

In our previous studies we have shown that CD4 T-cell death, in particular intrinsic apoptosis, is a major determinant of immune recovery [8, 17]. Therefore, we explored the association of unbalanced CD4 T-cell maturation with the rate of cell death in ex vivo cultures of fresh PBMC. Spontaneous CD4 T-cell death was unrelated to the frequency of CD4 T_{CM} or T_{TD} cells but showed a significant negative correlation with the frequency of CD4 TN and positive correlation with T_{TM} and T_{EM} cells (Figure 2a). Since the frequency of CD4 T_N and T_{TM} cells were strongly inversely correlated (data not shown), we addressed independent associations by using a model including data from all subsets. This model (Additional file 2: Table S1) confirmed the independent positive association of CD4 T-cell death with the frequency of T_{TM} CD4 T cells, clearly linking the higher presence of these cells with the increased cell death observed in immunodiscordant individuals.

However, these data do not discriminate whether TTM cell death rate is similar among treated HIV infected individuals or is higher in immunodiscordant patients. To experimentally confirm these possibilities, we addressed the analysis of the lifespan of T_N , T_{CM} , T_{TM} and T_{EM+TD} CD4 T-cells from immunocordant and immunodiscordant individuals. Since PBMC culture did not allow for a direct assessment of cell death occurring in different CD4 T-cell subsets due to phenotypic changes upon cell death [21], we chose a previously reported sorting strategy prior to ex vivo culture to approach this issue [21]. Although this strategy may induce some additional stress to sorted cells, it allowed for a consistent measure of subset cell death (Figure 2b). In this set of experiments, we observed that TN cells showed significantly lower cell death rates than those observed in purified memory cells in both groups (Figure 2c). No significant difference in TN cell death was observed between groups. However, memory cells in immunodiscordant individuals showed higher rates of death (Figure 2c). This was associated with higher sensitivity to cell death in TCM cells from immunodiscordant individuals ($p = 0.056$), and non-significant trends in TTM and T_{EM+TD} cells. Consistently, T_N cell death did not correlate with CD4 T-cell counts while a significant negative association ($p = 0.013$) was observed for T_{CM} and trends were noticed for T_{TM} and T_{EM+TD} cell death ($p = 0.067$ and 0.054 , respectively; Figure 2d). No significant correlations with senescence markers could be identified (data not shown).

To evaluate the impact of higher cell death rates on transition from T_{CM} to more advanced stages of CD4 T-cell maturation, we analyzed the ratio between T_{CM} and T_{TM} cells. This parameter was significantly lower in immunodiscordant subjects compared to immunocordant (irrespective of nadir values) or uninfected individuals (Figure 3a). Interestingly, T_{CM} to T_{TM} transition is not completely normalized in immunocordant individuals (Figure 3a), suggesting that this parameter could be useful to assess the quality of immune recovery. Reinforcing the relevance of the T_{CM}/T_{TM} ratio, a strong negative correlation was observed between this parameter and spontaneous CD4 T-cell death (Figure 3b). Overall, our data suggest that memory, but not T_N cells, are major contributors to the increased cell death observed in immunodiscordant individuals [17]. Higher sensitivity to death of memory cell populations, especially T_{CM} , may help to explain the inability of immunodiscordant individuals to recover proper T_{CM}/T_{TM} ratios.


CD57 expression and immune recovery

Besides T_{CM} or T_{TM} dysfunction, replicative immunosenescence of CD4 T cells has also been related to immune recovery [14]. In our cohort, analysis of the frequency of $CD28^-CD57^+$ CD4 T cells (Additional file 1: Figure S1) showed higher levels in immunodiscordant subjects than in both immunocordant subgroups, which in turn showed similar levels to HIV-uninfected individuals (Figure 4a). The analysis of CD57 expression in the different CD4 T-cell subsets showed that despite apparent full immune recovery, immunocordant individuals displayed higher CD57 expression than uninfected controls in all CD4 T-cell subsets. In addition, immunodiscordant individuals showed highly significant differences with concordant subjects particularly in T_N and T_{CM} cells (Figure 4b).

We also evaluated the relationship of CD57 expression with CD4 T-cell survival. No significant correlation was found between replicative senescence ($CD28^-CD57^+$ cells) and spontaneous CD4 T-cell death (Figure 4c). These data suggest that different mechanisms regulate the expression of CD57 in different CD4 T-cell subsets and show that replicative senescence is not coupled with the spontaneous CD4 T-cell death rate characteristics of immunodiscordant responses to HAART.


Analysis of the CD8 T-cell compartment

Major comorbidities in HIV infected individuals have been related to CD8 T-cell activation [1, 22] and recently to the CD4/CD8 T-cell ratio. Therefore, we also evaluated differentiation and immunosenescence in CD8 T cells. Elevated CD8 T-cell counts in HIV infected individuals analyzed in this study (Table 1) were the consequence of



an expansion of all subsets except CD8 T_{CM} cells (Figure 5a). Comparison of immunodiscordant and immunoconcordant subjects exclusively showed differences in the T_N subset, consistent with the widely described thymic dysfunction of these patients [6]. The percentage of CD8 T cells was not normalized in immunoconcordant individuals, which showed intermediate values between immunodiscordant and uninfected subjects (Table 1).

Among immunoconcordant and immunodiscordant individuals, T_{TM} , T_{EM} and T_{TD} CD8 T-cell subsets showed similar frequencies. However, a decreased size of the CD8 T_N subset was observed in immunodiscordant individuals, while a reduced T_{CM} subset was only observed in immunoconcordant subjects (Figure 5b). Replicative immunosenescence ($CD28^-CD57^+$ cells) was increased in both groups of HIV infected individuals




compared to uninfected controls; however, no differences between immunoconcordant and immunodiscordant individuals were observed. A deeper analysis of CD57 expression in the different CD8 T-cell subsets commonly showed a lower expression of CD57 in uninfected donors compared to HIV infected groups, with no differences between immunodiscordant and immunoconcordant individuals (Figure 5c). Again, we observed similar CD57 expression when this latter group was subdivided according to nadir values of CD4 T-cell counts. In summary, individuals displaying different CD4 T-cell recovery profiles seem to have similar CD8 T-cell compartments, except for the number of CD8 T_N cells.

Discussion

Immunodiscordant responses to HAART are associated with high risk of mortality and the occurrence of AIDS and non AIDS-related diseases [23–28]. Although blunted CD4 T-cell recovery has been associated with impaired thymic output [29], other factors affecting memory cell populations, such as activation, replicative immunosenescence [30], skewed T-cell maturation [15, 30, 31] and increased rates of CD4 T-cell death [8, 17], may contribute to persistently low CD4 T-cell counts.

The pivotal role of TN cells in CD4 T-cell repopulation and the lower size of this subset in immunodiscordant individuals has been universally reported [5, 10, 32, 33] and has been confirmed in our cohort using the CD31 marker (CD45RA⁺CD31⁺ percentage of CD4 T cells) [17]. However, the thymus hypothesis should not

overlook the role of memory cell populations, which may maintain T-cell homeostasis in thymectomized SIV-infected animals [11]. Immunodiscordant individuals show low proliferation of T_N cells [15], suggesting that T_N cells might be not required for CD4 T-cell repopulation (Additional file 3: Figure S2). In the absence of new CD4 TN cells, memory cells should be expanded by homeostatic mechanisms involving IL-7 or TCR stimulation to maintain CD4 T-cell numbers [34, 35]. Since CD4 T_{CM} cells are the main target of IL-7 [36] and IL-7 levels are increased in our cohort of immunodiscordant individuals [37], the inability to recover CD4 T cells, could also be related to the limited expansion capacity of memory cells. It could be argued that the CD4 TCM cell subset is under strong homeostatic pressure. Supporting this notion, high TCM proliferation have been reported in immunodiscordant individuals [15]. However, the limited size of this subset in immunodiscordant patients indicates that the proliferation of CD4 T_{CM} cells may be inefficient or may result in increased maturation towards T_{TM} or T_{EM} phenotypes. This could explain the relative accumulation of CD4 T cells in intermediate stages of differentiation, mainly TTM but also TEM cells, observed in our cohort of immunodiscordant individuals. Interestingly, our data show an association between the levels of CD4 T_{TM} cells and the rate of spontaneous CD4 T-cell death, although the analysis of cell death in sorted CD4 T-cell subsets in immunodiscordant and immunoconcordant individuals indicates that CD4 T_{CM} cell death is better correlated to CD4 T-cell recovery. Our data suggest that reduced T_{CM}




survival may be a second mechanism that adds to low thymic output in immunodiscordant patients.

An additional consequence of the high cell death level of T_{CM} and the trends detected in T_{TM} or T_{EM} cells in immunodiscordant patients is the modest accumulation of senescent CD4 T cells. Since the fate of CD4 T_{CM} and T_{TM} or T_{EM} could be cell death rather than further maturation (Additional file 3: Figure S2), this fact could explain the poor relationship between immunodiscordance and replicative senescence. Regarding senescence, our data on CD57 expression are consistent with previous reports [29], revealing higher levels of CD28⁻CD57⁺CD4 T cells in immunodiscordant individuals. Although some reports suggest that CD57⁺ CD4 T-cells undergo faster spontaneous apoptosis [38], the frequency of these cells is not associated with cell death rate in the CD4 T-cell compartment, suggesting that they are unlikely to drive increased cell death in immunodiscordant individuals.

Consistently, several authors suggest that CD28⁻CD57⁺ cells are long lived [39, 40]. Aside from replicative senescence, the expression of CD57 in other CD4 T-cell subsets, namely CD4 T_N cells, has been also associated with CD4 T-cell recovery [30]. Although we have also noticed this association, it remains unclear whether CD57⁺ cells in T_N and T_{CM} subsets are actually senescent [12], or reflect other T-cell functions or subsets such as Follicular helper CD4 T-cells (T_{Fh}), which express CD57 [41] and are recovered after HAART [42]. The increased expression of CD57 in CD4 T_N and T_{CM} cells from immunodiscordant individuals points to alterations in these subsets and suggests again a role for T_{CM} cells in CD4 T-cell recovery.

Our data point out that the level of destruction of the immune system, as measured by the nadir CD4 T-cell count, which has been widely associated with poor CD4 T-cell recovery [5], fail to fully identify



immunodiscordant responses. Indeed, some individuals with low nadir values are able to recover the CD4 T-cell compartment to the same extent as subjects with higher nadir values, at least in respect to maturation

phenotypes. Although this observation does not occlude the fact that starting HAART with low CD4 T-cell counts increases the risk of immunodiscordant responses, it suggests that not only the number but also the quality of CD4 T cells determine immune reconstitution. In this regard, it is tempting to speculate that the size of the TCM compartment prior to HAART may yield a better predictive value [16].

We have also shown that despite the impairment/failure of thymic output in both CD4 and CD8 T-cell subsets, only CD4 T cells show an unbalanced maturation profile. While minimal differences were observed among HIV infected groups in CD8 T cells. This result is surprising since CD8 T-cell senescence or activation has also been related to clinical progression in treated HIV infected individuals [43, 44]. Moreover, a recent analysis of HIV infected individuals failing to recover CD4/CD8 T-cell ratios after HAART also points to maturation and activation of CD8 T cells as major markers of poor immune recovery [45]. This apparent paradox could be related to the different pressures acting on CD4 and CD8 T cells or to the specific features of our cohort, which exclusively included long-term treated individuals (more than 2 years) without AIDS defining events. It remains possible that we have selected individuals with low risk of clinical progression, potentially minimizing differences in CD8 T cells. On the other hand, our selection criteria and classification cutoff, based exclusively in CD4 T-cell counts, allows for the direct comparison of the CD4 T-cell compartment between groups, avoiding the confounding effects of inflammatory and clinical status. In contrast, our analysis has some limitations. First, inclusion criteria might select immunodiscordant patients with a trend towards a lower treatment time (see Table 1). However, correcting maturation and senescence analyses for time on treatment did not modify the main conclusions of the study (not shown). A second limitation comes from the definition of immunodiscordant responses. We have used a cutoff value of 350 CD4 T cells/ μ L, while other authors define the limit in higher values. To assess the robustness of our analysis, a cutoff of 500 CD4 T cells/ μ L has been also used, yielding similar results (data not shown). Finally, another confounding factor relevant in the context of this study is CMV co-infection, a major driver of immunosenescence. However, all HIV infected individuals and 9 out of 11 uninfected individuals recruited were seropositive for CMV infection. Exclusion of CMV seronegative controls did not modify our analysis.

Conclusions

In summary, our data point to a major perturbation of the central memory CD4 T cells as a new prominent

characteristic of immunodiscordant responses to HAART. This perturbation adds to the increased accumulation of senescent cells and to the poor thymic output, and points to CD4 T_{CM} and T_{TM} cells as the major players of CD4 T-cell repopulation.

Additional files

Additional file 1: Figure S1. Gating strategy followed to identify T-cell maturation stages. **Panel A.** CD4 and CD8 T cells were gated and analyzed for the expression of CD27 and CD28. For CD4 T cells, terminally differentiated (T_{TD}) cells were defined as CD28⁻CD27⁻, while effector memory (T_{EM}) cells were CD28⁺CD27⁻. Double positive cells were further analyzed for CCR7 and CD45RA expression to identify naïve (T_N) cells (CCR7⁺CD45RA⁺), central memory (T_{CM}) cells (CCR7⁺CD45RA⁻) and transitional Memory (T_{TM}) cells (CCR7⁻CD45RA⁻). **Panel B.** For CD8 T cells, the general strategy was similar, unless for the definition of T_{EM} cells that was CD27⁺CD28⁻ cells. The expression of CD57 was analyzed in the whole CD4 or CD8 T-cell population (lower left dot plot in each panel) to define replicative senescence (CD28⁻CD57⁺ cells). In addition the expression of CD57 in each subset was also evaluated (lower dot plots in each panels).

Additional file 2: Table S1. A. The multivariate relationship between T_N , T_{CM} , T_{TM} , T_{EM} , T_{TD} and Total Death were determined by multiple linear regression. **B.** Linear regression model shows no statistical differences between group of patients (Discordant/Concordant) in the association of Total Death with T_{TM} .

Additional file 3: Figure S2. A model for CD4 homeostasis in Immunodiscordant individuals. Immunoconcordant individuals show a full recovery of CD4 T-cell counts with a high representation of naïve cells, and balanced frequencies of different memory subsets (upper panels). The profile of CD4 T-cell maturation in immunoconcordant individuals (green line) overlaps with that of HIV uninfected individuals (blue line in upper right plot) in which CCR7⁻ CD4 T cells show increased turnover and short live [22]. Conversely, immunodiscordant individuals (lower panels) show a reduced naïve subset with no signs of altered turnover [14] that limits the generation of new memory cells. Among memory subsets, central memory cells are also reduced and subjected to homeostatic pressure to generate new cells [14] increasing T_{CM} → T_{TM} transition (and lowering T_{CM}/T_{TM} ratios) and increasing T_{CM} cell death. T_{TM} and T_{EM} cells emerging from T_{CM} cells also show higher sensitivity to cell death (Figure 2D), explaining the lack of accumulation of terminally differentiated cells in these subjects. This scenario results in a shifted profile of CD4 T-cell maturation (red line) compared to healthy individuals (blue line in lower right plot).

Authors' contributions

MM, MC, EG and JC performed flow cytometry data acquisition, analysis, and interpretation, and revised the manuscript; DO performed statistical data analysis; JP and CC contributed to data analysis and interpretation, and revised the manuscript; EN and BC contributed to study design, clinical follow up and revised the manuscript; JB contributed to study design and data interpretation and wrote the manuscript. All authors read and approved the final manuscript.

Author details

¹ Institut de Recerca de la Sida IrsiCaixa-HIVACAT, Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Universitat Autònoma de Barcelona, 08916 Badalona, Spain. ² Fundació Lluita contra la SIDA, Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, 08916 Badalona, Spain. ³ Universitat de Vic-Central de Catalunya, UVIC-UCC, 08500 Vic, Spain. ⁴ Present Address: Department of Pathology, University of California San Diego, La Jolla, CA 92093, USA.

Acknowledgements

We are grateful to Leontina Banica (National Institute for Infectious Diseases, Bucharest) for critical reading of the manuscript, to technical staff of IrsiCaixa for sample processing and clinical staff of Fundació Lluita contra la Sida for clinical monitoring. This work was supported by the HIVACAT Program and the

Spanish AIDS network 'Red Temática Cooperativa de Investigación en SIDA RD12/0017/0002 project as part of the Plan Nacional R + D + I and cofunded by ISCIII- Subdirección General de Evaluación y el Fondo Europeo de Desarrollo Regional (FEDER). J.B. and C. C. are researchers from Fundació Institut de Recerca en Ciències de la Salut Germans Trias i Pujol supported by the ISCIII and the Health Department of the Catalan Government (Generalitat de Catalunya). J.C. is supported by a 'Sara Borrell' grant from the Spanish Health Institute 'ISCIII'.

Compliance with ethical guidelines

Competing interests

All authors declare that they have no competing interests related to this work. Data have been partly presented at CROI meeting 2013.

Received: 9 April 2015 Accepted: 9 July 2015

Published online: 17 July 2015

References

- Deeks SG (2011) HIV infection, inflammation, immunosenescence, and aging. *Annu Rev Med* 62:141–155
- Blanco J, Barretina J, Cabrera C, Gutiérrez A, Clotet B, Esté JA (2001) CD4(+) and CD8(-) T cell death during human immunodeficiency virus infection in vitro. *Virology* 285:356–365
- Doitsh G, Cavrois M, Lassen KG, Zepeda O, Yang Z, Santiago ML et al (2010) Abortive HIV infection mediates CD4 T cell depletion and inflammation in human lymphoid tissue. *Cell* 143:789–801
- Breton G, Chomont N, Takata H, Fromentin R, Ahlers J, Filali-Mouhim A et al (2013) Programmed death-1 is a marker for abnormal distribution of naïve/memory T cell subsets in HIV-1 infection. *J Immunol* 191:2194–2204
- Gazzola L, Tincati C, Bellistri GM, Monforte A, Marchetti G (2009) The absence of CD4⁺ T cell count recovery despite receipt of virologically suppressive highly active antiretroviral therapy: clinical risk, immunological gaps, and therapeutic options. *Clin Infect Dis* 48:328–337
- Massanella M, Negredo E, Clotet B, Blanco J (2013) Immunodiscordant responses to HAART—mechanisms and consequences. *Expert Rev Clin Immunol* 9:1135–1149
- Corbeau P, Reynes J (2011) Immune reconstitution under antiretroviral therapy: the new challenge in HIV-1 infection. *Blood* 117:5582–5590
- Negredo E, Massanella M, Puig J, Pérez-Alvarez N, Gallego-Escuredo JM, Villarroya J et al (2010) Nadir CD4 T cell count as predictor and high CD4 T cell intrinsic apoptosis as final mechanism of poor CD4 T cell recovery in virologically suppressed HIV-infected patients: clinical implications. *Clin Infect Dis* 50:1300–1308
- Marchetti G, Gori A, Casabianca A, Magnani M, Franzetti F, Clerici M et al (2006) Comparative analysis of T-cell turnover and homeostatic parameters in HIV-infected patients with discordant immune-virological responses to HAART. *AIDS* 20:1727–1736
- Ruiz-Mateos E, Rubio A, Vallejo A, De la Rosa R, Sanchez-Quijano A, Lissen E et al (2004) Thymic volume is associated independently with the magnitude of short- and long-term repopulation of CD4⁺ T cells in HIV-infected adults after highly active antiretroviral therapy (HAART). *Clin Exp Immunol* 136:501–506
- Okoye AA, Rohankhedkar M, Abana C, Pattann A, Reyes M, Pexton C et al (2012) Naïve T cells are dispensable for memory CD4⁺ T cell homeostasis in progressive simian immunodeficiency virus infection. *J Exp Med* 209:641–651
- Brenchley JM, Karandikar NJ, Betts MR, Ambrozak DR, Hill BJ, Crotty LE et al (2003) Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8⁺ T cells. *Blood* 101:2711–2720
- Fernandez S, Price P, McKinnon EJ, Nolan RC, French MA (2006) Low CD4⁺ T-cell counts in HIV patients receiving effective antiretroviral therapy are associated with CD4⁺ T-cell activation and senescence but not with lower effector memory T-cell function. *Clin Immunol* 120:163–170
- Méndez-Lagares G, Díaz L, Correa-Rocha R, León Leal JA, Ferrando-Martínez S, Ruiz-Mateos E et al (2013) Specific patterns of CD4-associated

- immunosenescence in vertically HIV-infected subjects. *Clin Microbiol Infect* 19:558–565
15. Lederman MM, Calabrese L, Funderburg NT, Clagett B, Medvik K, Bonilla H et al (2011) Immunologic failure despite suppressive antiretroviral therapy is related to activation and turnover of memory CD4 cells. *J Infect Dis* 204:1217–1226
 16. Okoye A, Meier-Schellersheim M, Brechley JM, Hagen SI, Walker JM, Rohankhedkar M et al (2007) Progressive CD4⁺ central memory T cell decline results in CD4⁺ effector memory insufficiency and overt disease in chronic SIV infection. *J Exp Med* 204:2171–2185
 17. Massanella M, Negro E, Pérez-Alvarez N, Puig J, Ruiz-Hernández R, Bofill M et al (2010) CD4 T-cell hyperactivation and susceptibility to cell death determine poor CD4 T-cell recovery during suppressive HAART. *AIDS* 24:959–968
 18. Curriu M, Carrillo J, Massanella M, Rigau J, Alegre J, Puig J et al (2013) Screening NK-, B- and T-cell phenotype and function in patients suffering from Chronic Fatigue Syndrome. *J Transl Med* 11:68
 19. Massanella M, Curriu M, Carrillo J, Gómez E, Puig J, Navarro J et al (2013) Assessing main death pathways in T lymphocytes from HIV infected individuals. *Cytometry A* 83:648–658
 20. R Core Team (2012) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. <http://www.R-project.org>
 21. Chirullo B, Sgarbanti R, Limongi D, Shytaj IL, Alvarez D, Das B et al (2013) A candidate anti-HIV reservoir compound, auranofin, exerts a selective 'anti-memory' effect by exploiting the baseline oxidative status of lymphocytes. *Cell Death Dis* 4:e944
 22. Deeks SG (2009) Immune dysfunction, inflammation, and accelerated aging in patients on antiretroviral therapy. *Top HIV Med* 17:118–123
 23. Kaufmann GR, Furrer H, Ledergerber B, Perrin L, Opravil M, Vernazza P et al (2005) Swiss HIV Cohort Study: Characteristics, determinants, and clinical relevance of CD4 T cell recovery to <500 cells/microL in HIV type 1-infected individuals receiving potent antiretroviral therapy. *Clin Infect Dis* 41:361–372
 24. Gutiérrez F, Padilla S, Masiá M, Iribarren JA, Moreno S, Viciano P et al (2008) Patients' characteristics and clinical implications of suboptimal CD4 T-cell gains after 1 year of successful antiretroviral therapy. *Curr HIV Res* 6:100–107
 25. Piketty C, Weiss L, Thomas F, Mohamed AS, Belec L, Kazatchkine MD (2001) Long-term clinical outcome of human immunodeficiency virus-infected patients with discordant immunologic and virologic responses to a protease inhibitor-containing regimen. *J Infect Dis* 183:1328–1335
 26. Young J, Psychogiou M, Meyer L, Ayayi S, Grabar S, Raffi F et al (2012) CD4 cell count and the risk of AIDS or death in HIV-infected adults on combination antiretroviral therapy with a suppressed viral load: a longitudinal cohort study from COHERE. *PLoS Med* 9:e1001194
 27. Tan R, Westfall AO, Willig JH, Mugavero MJ, Saag MS, Kaslow RA et al (2008) Clinical outcome of HIV-infected antiretroviral-naive patients with discordant immunologic and virologic responses to highly active antiretroviral therapy. *J Acquir Immune Defic Syndr* 47:553–558
 28. Moore DM, Hogg RS, Chan K, Tyndall M, Yip B, Montaner JSG (2006) Disease progression in patients with virological suppression in response to HAART is associated with the degree of immunological response. *AIDS* 20:371–377
 29. Appay V, Fastenackels S, Katlama C, Ait-Mohand H, Schneider L, Guihot A et al (2011) Old age and anti-cytomegalovirus immunity are associated with altered T-cell reconstitution in HIV-1-infected patients. *AIDS* 25:1813–1822
 30. Méndez-Lagares G, García-Pergañeda A, del Mardel Pozo-Balado M, Genebat M, Ruiz-Mateos E, García García M et al (2012) Differential alterations of the CD4 and CD8 T cell subsets in HIV-infected patients on highly active antiretroviral therapy with low CD4 T cell restoration. *J Antimicrob Chemother* 67:1228–1237
 31. Marchetti G, Gazzola L, Trabattini D, Bai F, Ancona G, Ferraris L et al (2010) Skewed T-cell maturation and function in HIV-infected patients failing CD4⁺ recovery upon long-term virologically suppressive HAART. *AIDS* 24:1455–1460
 32. Solomon A, Cameron PU, Bailey M, Dunne AL, Crowe SM, Hoy JF et al (2003) Immunological and virological failure after antiretroviral therapy is associated with enhanced peripheral and thymic pathogenicity. *J Infect Dis* 187:1915–1923
 33. Smith KY, Valdez H, Landay A, Spritzler J, Kessler HA, Connick E et al (2000) Thymic size and lymphocyte restoration in patients with human immunodeficiency virus infection after 48 weeks of zidovudine, lamivudine, and zalcitabine therapy. *J Infect Dis* 181:141–147
 34. Riou C, Yassine-Diab B, Van Grevenynghe J, Somogyi R, Grellier LD, Gagnon D et al (2007) Convergence of TCR and cytokine signaling leads to FOXO3a phosphorylation and drives the survival of CD4⁺ central memory T cells. *J Exp Med* 204:79–91
 35. Catalfamo M, Di Mascio M, Hu Z, Srinivasula S, Thaker V, Adelsberger J et al (2008) HIV infection-associated immune activation occurs by two distinct pathways that differentially affect CD4 and CD8 T cells. *Proc Natl Acad Sci USA* 105:19851–19856
 36. Sereti I, Dunham RM, Spritzler J, Aga E, Proschan MA, Medvik K et al (2009) IL-7 administration drives T cell-cycle entry and expansion in HIV-1 infection. *Blood* 113:6304–6314
 37. Massanella M, Ouchi D, Marfil S, Llibre JM, Puertas MC, Buzón MJ et al (2014) Different Plasma Markers of Inflammation Are Influenced by Immune Recovery and cART Composition or Intensification in Treated HIV Infected Individuals. *PLoS One* 9:e114142
 38. Palmer BE, Blyveis N, Fontenot AP, Wilson CC (2005) Functional and phenotypic characterization of CD57⁺CD4⁺T cells and their association with HIV-1-induced T cell dysfunction. *J Immunol* 175:8415–8423
 39. Spaulding C, Guo W, Effros RB (1999) Resistance to apoptosis in human CD8⁺ T cells that reach replicative senescence after multiple rounds of antigen-specific proliferation. *Exp Gerontol* 34:633–644
 40. Gupta S (2005) Molecular mechanisms of apoptosis in the cells of the immune system in human aging. *Immunol Rev* 205:114–129
 41. Bossaller L, Burger J, Draeger R, Grimbacher B, Knöth R, Plebani A et al (2006) ICOS deficiency is associated with a severe reduction of CXCR5⁺CD4 germinal center Th cells. *J Immunol* 177:4927–4932
 42. Boswell KL, Paris R, Boritz E, Ambrozak D, Yamamoto T, Darko S et al (2014) Loss of circulating CD4 T cells with B cell helper function during chronic HIV infection. *PLoS Pathog* 10:e1003853
 43. Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ et al (2011) T cell activation predicts carotid artery stiffness among HIV-infected women. *Atherosclerosis* 217:207–213
 44. Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ et al (2011) T cell activation and senescence predict subclinical carotid artery disease in HIV-infected women. *J Infect Dis* 203:452–463
 45. Serrano-Villar S, Sainz T, Lee SA, Hunt PW, Sinclair E, Shacklett BL et al (2014) HIV-Infected Individuals with Low CD4/CD8 Ratio despite Effective Antiretroviral Therapy Exhibit Altered T Cell Subsets, Heightened CD8⁺ T Cell Activation, and Increased Risk of Non-AIDS Morbidity and Mortality. *PLoS Pathog* 10:e1004078

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

